





## ***Prevalence of Hypothyroidism and Pregnancy Outcomes in Women Referred to Ayatollah Mousavi Hospital in Zanjan in 2018-2019***

Bayat F<sup>1</sup>, Rezaee Moradali M<sup>2</sup>, Roozbahani S<sup>2</sup>, Towfighi Sh<sup>3</sup>, Shami M<sup>4\*</sup>

<sup>1</sup>Ph.D student, Instructor, Department of Midwifery, School of Nursing and Midwifery, Zanjan University of Medical Sciences, Zanjan, Iran

<sup>2</sup>Ph.D student of reproductive health, Student Research Committee, Midwifery and Reproductive Health Research Center, School of Nursing and Midwifery, Shahid Beheshti University of Medical Sciences, Tehran, Iran

<sup>3</sup>Assistant Professor of Obstetrics and Gynecology, Department of Obstetrics and Gynecology, School of Medicine, Ayatollah Mousavi Hospital, Zanjan University of Medical Sciences

<sup>4</sup>master student, Tehran university of medical sciences, school of nursing and midwifery, Tehran, Iran

**\*Corresponding Author:** Department of Midwifery and Reproductive Health, School of Nursing and Midwifery, Tehran University of Medical Sciences, Tehran, Iran

**Tel:** 0098- 9199549600

**Email:** maryamshami1870@gmail.com

**Received:** 4 Oct 2020

**Accepted:** 23 Nov 2020

### **Abstract**

**Background:** Hypothyroidism is common during pregnancy. This disorder is associated with adverse maternal and neonatal outcomes.

**Objectives:** The study of the prevalence and different consequences of hypothyroidism in pregnancy in each region is the special importance in determining prevention and management strategies. Therefore, the aim of this study was to determine the prevalence of hypothyroidism and pregnancy outcomes in women referred to Ayatollah Mousavi Hospital in Zanjan.

**Methods:** This cross-sectional-analytical study was performed on all pregnant women who gave birth in Ayatollah Mousavi Hospital in Zanjan in 2018-2019. Data were collected through census and using Chi-square, Fisher's exact and independent t-tests and with SPSS software version 22, pregnancy outcomes between the two groups were calculated.

**Results:** The prevalence of hypothyroidism in pregnant women in the present study was estimated at 8.59%. Hypothyroidism was found to double the chance of gestational diabetes ( $p<0.02$ ) (CI=1.1-3.8). The rate of cesarean delivery was significantly higher in patients ( $p<0.001$ ). The rate of fetal heart rate drop, placental abruption, and placental adhesion was higher in the affected group, but this difference was not significant ( $p>0.05$ ). There was no significant difference between the two groups in terms of preeclampsia, preterm delivery, first minute Apgar score, height, weight and head circumference of the newborn ( $p>0.05$ ).

**Conclusion:** Considering the adverse maternal and neonatal outcomes of hypothyroidism, it seems that more clinical studies in this field are needed to manage and evaluate the outcomes in pregnant women with hypothyroidism.

**Keywords:** *prevalence, hypothyroidism, pregnancy outcomes, neonatal outcomes*

### **Introduction**

During the pregnancy the thyroid undergoes physiology and structure changes, and the fetus is reliant on maternal thyroxine, so the mother's thyroid gland is more prone to hypothyroidism during pregnancy [1].

Various factors influence the function of the thyroid during pregnancy such as adequate iodine intake, habits and diet, environmental factors, and genetic predisposition [2]. Maternal age, number

of pregnancies, living area, history of infertility, history of abortion, using birth control pills, presence of ovarian cysts, smoking, high body mass index, thyroid autoimmune diseases, thyroid gland destruction with radioactive iodine for the treatment of hyperthyroidism and a history of thyroid disorders [2-4].

Several studies have mentioned the association between iodine and hypothyroidism, so that in areas of iodine deficiency, the size of the thyroid

increases by 20-40% and the prevalence of hypothyroidism is higher. More than one third of the world's population suffer from iodine deficiency. It's believed that the prevalence of goiter is higher in mountainous areas due to iodine deficiency, and since Zanjan province is also geographically located in a mountainous region, the prevalence of hypothyroidism is on the rise. On the other hand, pregnancy is associated with an increased need for daily iodine intake by 50%, mainly due to increased iodine excretion through the kidneys [1,5,6].

The most common thyroid dysfunction in pregnancy is hypothyroidism [7], nearly 2 to 12 in 1000 pregnancies [1]. According to the literature, the prevalence of this disorder based on the criteria of the American Thyroid Association (ATA), in western countries ranges from 0.5 to 12.3% [8,9]. Meanwhile, its prevalence is higher in Asian countries (13.13-14%) [10,11]. In Iran, the prevalence of hypothyroidism varies according to geography and the amount of iodine intake [6]. Nazarpour et al. have reported a prevalence of 18.9 and 50% for subclinical and overt hypothyroidism in pregnancy in the city of Tehran, respectively [3]. Also, its prevalence is estimated to be 21% and 10.15% in Mashhad (northeast of Iran) and Kerman (east of Iran), respectively [12]. In Hamedan, which is one of the neighboring provinces of Zanjan, the prevalence of hypothyroidism during pregnancy in 2016-2017 was estimated to be 27.7% [13].

Hypothyroidism during pregnancy causes negative consequences for both mother and fetus. Hypothyroidism has two main types of overt and subclinical. Severe hypothyroidism is uncommon during pregnancy; mainly because severe hypothyroidism is less common during pregnancy due to infertility and recurrent miscarriage [1,7].

Ajmani et al. reported that those who suffer from hypothyroidism experience increased incidence of preeclampsia and placental abruption, but Mannisto studies reported no association between hypothyroidism and preeclampsia and preterm delivery [14-16]. Turunen et al. reported that maternal hypothyroidism was associated with gestational diabetes, hypertension in pregnancy and severe preeclampsia, cesarean section (C-section), preterm delivery, large infant relative to gestational age, major congenital anomalies, and neonatal hospitalization in the intensive care unit

(NICU). While Ezzeddine et al. did not find any association between maternal hypothyroidism and neonatal complications such as low Apgar score, low birth weight, and intrauterine death [2,17].

Based on what was mentioned before, there is a discrepancy in the findings and more studies are needed to evaluate the consequences of pregnancy in people with hypothyroidism. Also, according to the available knowledge, no studies investigated the prevalence of hypothyroidism in Zanjan. Therefore, this study was performed to estimate the prevalence of hypothyroidism in women referred to Ayatollah Mousavi Hospital in Zanjan. Besides pregnancy and neonatal related outcomes are also investigated.

## Methods

This cross-sectional analytical study was performed on all pregnant women who had given birth in Ayatollah Mousavi Hospital of Zanjan from September 2018 to March 2019. The study was approved by the ethics committee of Zanjan University of Medical Sciences (code: A-12-1203-1) and ethical code (IR.ZUMS.REC.1397.166). The inclusion criteria were as follows gestation age of 22 weeks or higher, Iranian nationality, and singleton pregnancy. Exclusion criteria were having an underlying disease.

In the first phase of the study, a census was performed to identify all women who gave birth in the Hospital to determine the prevalence of hypothyroidism. Then, those with hypothyroidism were compared with others gestational diabetes and gestational disorders (Table 2). In the second phase, to compare the pregnancy outcomes in subjects with and without hypothyroidism, 115 mothers without hypothyroidism were selected by systematic random sampling. Therefore, 115 mothers with hypothyroidism and 115 mothers without hypothyroidism were studied, with a total sample size of 230.

Data were collected with a research checklist (maternal age, gestational age, neonatal sex, neonatal resuscitation need, all fetal heart rate deceleration, type of delivery, and neonatal outcome). Diagnosis of hypothyroidism was confirmed based on the medical history of patients, who was taken by a resident of obstetrics and gynecology; and evidence of hypothyroidism were according to the Ministry of Health, TSH

above 3.9 in the first trimester or TSH above 4.1 in the second and third trimesters. Data were analyzed using Chi-square, Fisher's exact test, and independent t-test by SPSS version 22. Statistical significance was considered when  $p\text{-value} < 0.05$ .

## Results

Of 1338 deliveries in the hospital, 115 mothers with hypothyroidism were identified, an estimated prevalence of 8.59%.

According to the Chi-square and Fisher's exact test, there was no statistically significant difference between the two groups concerning maternal age, place of residence, level of education, and history of abortion (Table 1). The number of primiparous women with hypothyroidism was higher than primiparous women in the control group, but this difference was not statistically significant ( $p > 0.05$ ).

**Table 1: Demographic and midwifery characteristics of participants**

variables	hypothyroidism Group Number (percent)	Control Group Number (percent)	P value
Age of Mother (Year)	30±2.6	8.29±6.6	0.822 <sup>a</sup>
Area of Residence	City	61 (53)	0.182 <sup>b</sup>
	Village	54 (47)	
Educational Status	illiterate	6 (2.5)	0.387 <sup>b</sup>
	Secondary	93 (9.80)	
	University	16 (9.13)	
Parity	Nulliparous	27 (5.23)	0.059 <sup>b</sup>
	Multiparous	88 (5.76)	
Abortion Record	Yes	19 (5.16)	0.076 <sup>b</sup>
	No	96 (5.83)	

Independent t-test <sup>a</sup>

Chi-square or Fisher exact test <sup>b</sup>

In this study, hypothyroidism was significantly associated with gestational diabetes ( $P < 0.02$ ), so that those with hypothyroidism were at increased risk of developing gestational diabetes by two

times (CI=1.1-3.8). But no significant association was found between hypothyroidism and hypertension disorders ( $P > 0.05$ ) (Table 2).

**Table 2: Comparison of gestational diabetes and hypertensive disorders in study participants**

variables	Hypothyroidism Yes (115 mothers)	Hypothyroidism No (1223 mothers)	P value <sup>a</sup>	OR (CI)
Disease	hypertensive disorders	75 (6.1)	0.720	(0.53-2.43) 1.14
	gestational diabetes	71 (5.8)	0.020*	(1.3-1.8) 2.06

Chi-square <sup>a</sup>

In this study, the rate of C-section in the affected group was significantly higher ( $P < 0.001$ ) so that hypothyroidism was associated risk of C-section by 2.5 times. Fetal meconium excretion was more common among those with normal pregnancy

( $P < 0.03$ ). Fetal heart rate deceleration, placental abruption, and placental adhesion were higher in mothers with hypothyroidism, but this difference was not statistically significant ( $P > 0.05$ ) (Table 3).

**Table 3: Comparison of pregnancy outcome between two group**

variables		Hypothyroidism Group Number (percent)	Control Group Number (percent)	P Value	OR (CI)
<b>Mode of Delivery</b>	Natural	39 (33.9)	65 (56.5)	0.001 <sup>a</sup>	(1.4-4.3) 2.5
	Cesarean	76 (66.1)	50 (43.5)		
<b>Meconium</b>	Yes	1 (0.9)	7 (6.1)	0.033 <sup>b*</sup>	(0.1-1.1) 0.13
	No	114 (99.1)	108 (93.9)		
<b>Fetal heart rate deceleration</b>	Yes	19 (16.5)	10 (8.7)	0.076 <sup>a</sup>	(0.4-92.69) 2.07
	No	96 (83.5)	105 (91.3)		
<b>Placental abruption</b>	Yes	5 (4.3)	1 (0.9)	0.106 <sup>b</sup>	-
	No	110 (95.7)	114 (99.1)		
<b>Placental adhesion</b>	Yes	4 (3.5)	0 (0)	0.061 <sup>b</sup>	-
	No	111 (96.5)	115 (100)		

a: chi-square

b: Fisher's exact test

\* significant

According to the independent t-test, neonatal outcomes were not significantly different between

the two groups (Table 4).

**Table 4: Comparison of neonatal outcomes between two groups**

variables	Hypothyroidism Group	Control Group	P-Value <sup>a</sup>
<b>Age of pregnancy</b>	37.2±5.7	37.2±2.9	0.465
<b>Birth weight</b>	3079.698±39.552	2987.681±72.09	0.313
<b>Height</b>	50.4±2.3	49.3±8.79	0.562
<b>Head circumference</b>	34.2±6	34.2±5.9	0.201
<b>Apgar first minute</b>	8.1±2.8	8.1±2.2	0.762

**Independent-Sample T test<sup>a</sup>**

## Discussion

In the present study, the prevalence of hypothyroidism in pregnant women referred to Ayatollah Mousavi Hospital in Zanjan was 8.59%, which is consistent with some of the previously conducted studies. For example, the prevalence of this disorder in the cities of Hamadan, Kerman, and Mashhad has been estimated at 1.2%, 10.15%, and 21%, respectively [12,13,18]. A recently conducted study in the city of Tehran reported a prevalence of 9.9% for hypothyroidism during pregnancy [5]. The prevalence of this disorder according to the criteria of the American Thyroid Association in Western countries is estimated at 0.3-2.79% [8,9,19]. In contrast, based on recent studies conducted in Asian countries, the prevalence of hypothyroidism is higher in these countries, ranging from 13.13 to 14% [10,11]. It seems that

geography also contributes to the prevalence of hypothyroidism. Besides, sample size and study protocol are also important factors in estimating the prevalence of hypothyroidism.

Based on the findings, the prevalence of hypothyroidism was higher in nulliparous women than their multiparous counterparts but this difference was not significant ( $p>0.05$ ). Cleary-Goldman et al. also, reported that the prevalence of hypothyroidism was higher in nulliparous women than multiparous women [20]. Consistent with the present study, Nazarpour et al. [3] and Dulek et al. [9] also reported no association between parity and hypothyroidism. It seems that their sample size has influenced the results.

Similar to Kiran et al., this study demonstrated that hypothyroidism was not associated with a history of abortion ( $p=0.07$ ) [11] which is in contrast to the study by Ajmani et al., Shinohara

et al., and Cakmak et al., who reported that hypothyroidism was associated with abortion and that the prevalence of abortion was higher in people with hypothyroidism [15,21,22]. This difference can be attributed to the fact that whether patients were receiving treatment or not. Ajmani et al. reported a higher prevalence of preeclampsia among women with hypothyroidism so that these women were twice as likely as their healthy counterparts at risk of developing preeclampsia [15]. Kiran et al. also reported that the incidence of preeclampsia increases following hypothyroidism [11]. These findings are in contrast to the results of the present study, which showed no significant association between hypothyroidism and hypertensive disorders. As in our study, Ezzeddine et al. also found no significant association between hypothyroidism and hypertension during pregnancy [2]. This discrepancy can be attributed to the fact that in the present study, women's hypothyroidism was under control and treatment.

Chen et al. reported no significant association between hypothyroidism and gestational diabetes [23]. But the results of our study showed that people with hypothyroidism are at increased risk of developing gestational diabetes (about two times). Stohl et al. also emphasized that hypothyroidism is strongly associated with gestational diabetes and recommended that screening for diabetes should begin earlier in women with thyroid dysfunction [24]. The observed difference can be attributed to factors such as the methodology of the study and criteria used for diagnosing hypothyroidism.

Based on the findings, the rate of C-section in women with hypothyroidism was 2.5 times higher than in healthy women. This increase may be secondary to an increase in gestational diabetes in women with hypothyroidism. Stohl et al. showed that the prevalence of gestational diabetes is higher in women with hypothyroidism than those with hyperthyroidism [24]. Cakmak et al. also reported similar results. According to their study, the risk of C-section was 1.2 times higher in women with hypothyroidism. They argued that the observed increase in the rate of C-section can be attributed to the enhanced rate of pregnancy related outcomes such as high blood pressure, diabetes, and low birth weight [21]. The findings of the present study are consistent with the results

of Kiran et al. who reported that TSH levels  $> 2.5$  were associated with an increased rate of C-section [11]. Ajmani et al. argued that the cause of increased cesarean section was increased fetal distress [15]. In our study, the rate of heart rate drop was higher in the hypothyroidism group. But this difference was not statistically significant and could be related to the small sample size of the present study. In contrast, Dulek et al. reported in their study that there was no association between C-section and thyroid disorders during pregnancy [9]. Lee et al. found no association between high levels of TSH ( $TSH > 4$ ) and C-section [25]. These differences may be related to the study method, sample size, and the occurrence of other consequences.

No significant difference was found between the two groups concerning the neonatal outcomes (gestational age at birth, weight, height, head circumference, and first minute Apgar score). Similar to the results of the present study, Ezzeddine et al. reported no association between hypothyroidism and neonatal outcomes such as neonatal weight, 1 and 5 minutes Apgar scores less than 7, and intrauterine death [2]. Also Dulek et al. found no association between hypothyroidism and birth weight and Apgar scores at 1 and 5 minutes [9]. Cakmak et al. did not report any association between hypothyroidism and other neonatal outcomes such as preterm delivery, Apgar scores less than 7, low birth weight, and premature rupture of membranes, except for an increased risk of NICU (Neonatal Intensive Care Unit) admission [21]. Luisi et al. found no association between hypothyroidism and neonatal outcomes (low birth weight, head circumference, and height) [26]. In contrast, Lee et al. reported an increased risk of preterm infancy and respiratory distress syndrome for those with increased levels of TSH. Low birth weight and NICU admission were not associated with hypothyroidism [25]. Because subjects of the present study were receiving treatment their TSH were in the normal range, hence this difference can be justified.

### Conclusion

As women with hypothyroidism are at increased risk of hypothyroidism, it seems that more attention should be paid to screening pregnant women. Besides, to prevent the consequences of

hypothyroidism during pregnancy a series of programs should be developed. Further clinical studies are recommended to investigate other consequences of this disorder. The present research was a cross-sectional study and the subjects were treated with levothyroxine but the dose of the drug was not known. Using other study designs (e.g. cohort or RCT) would provide useful information. Moreover, as various studies have reported different rates of prevalence for hypothyroidism, it seems that newer and more effective criteria should be found for screening these people.

### Acknowledgements

This study was confirmed by the ethics committee of Zanjan University of Medical Sciences (code: A-12-1203-1) and ethical code (IR.ZUMS.REC.1397.166). We would like to thank all the participants who helped us with this project.

### Conflict of interest

The authors declare no conflict of interest.

### References

1. Dashe JS, Bloom SL, Spong CY, Hoffman BL. Williams obstetrics. 25<sup>th</sup> ed. New York: McGraw Hill Professional; 2018.
2. Ezzeddine D, Ezzeddine D, Hamadi C, et al. Prevalence and correlation of hypothyroidism with pregnancy outcomes among lebanese women. *J Endocr Soc*. 2017; 1(5): 415- 22.
3. Nazarpour S, Tehrani FR, Rahmati M, et al. Validation of Billewicz scoring system for detection of overt hypothyroidism during pregnancy. *Int J Endocrinol Metab*. 2018; 16(3).
4. Momtazan M, Mohammadi MJ, Tabahfar R, et al. Risk factors accelerating hypothyroidism in pregnant women referred to health centers in Abadan, Iran. *Data Brief*. 2017; 14: 15- 19.
5. Etemadi A, Amouzegar A, Mehran L, et al. Isolated Hypothyroxinemia in Iranian Pregnant Women, the Role of Iodine Deficiency: A Population-Based Cross-Sectional Study. *Thyroid*. 2020; 30(2): 262- 9.
6. Azizi F, Delshad MH, Amouzegar A, Mehran L, Mirmiran P. Prevention of iodine-deficiency induced goiter and brain damage in the islamic republic of Iran. *Iran J Endocrinol Metab*. 2014; 15(5): 411- 29.
7. Springer D, Jiskra J, Limanova Z, Zima T, Potlukova E. Thyroid in pregnancy: From physiology to screening. *Crit Rev in Clin Lab Sci*. 2017; 54(2): 102- 16.
8. López-Muñoz E, Ibarra-Avalos JA, Chan-Verdugo RG, Mateos-Sánchez L, Sánchez-Rodríguez O. Prevalence of hypothyroidism during pregnancy in a highly specialised referral centre in Mexico. *J Obstet Gynaecol*. 2016; 36(8): 1069- 75.
9. Dulek H, Vural F, Aka N, Zengin S. The prevalence of thyroid dysfunction and its relationship with perinatal outcomes in pregnant women in the third trimester. *North clin Istanbul*. 2019; 6(3): 267- 72.
10. Dhanwal DK, Bajaj S, Rajput R, et al. Prevalence of hypothyroidism in pregnancy: An epidemiological study from 11 cities in 9 states of India. *Indian J Endocrinol Metab*. 2016; 20(3): 387- 90.
11. Kiran Z, Sheikh A, Malik S, et al. Maternal characteristics and outcomes affected by hypothyroidism during pregnancy (maternal hypothyroidism on pregnancy outcomes, MHPO-1). *BMC Pregnancy childbirth*. 2019; 19(1): 476.
12. Lotfalizadeh M, Ghomian N, Mohammad Nezhad M. Prevalence and Complications of Hypothyroidism in Pregnancy. *Iran J Obstet, Gynecol Infertil*. 2017; 20(8): 1- 5. [In Persian]
13. Borzouei Sh, Goodarzi MT, Biglari M, Nazari F, Shivapour Z. The Prevalence of Thyroid Disorders in Pregnant Women of Hamadan. *Avicenna J Nurs Midwifery Care*. 2019; 27(1): 11- 7. [In Persian]
14. Ajmani SN, Aggarwal D, Bhatia P, Sharma M, Sarabhai V, Paul M. Prevalence of overt and subclinical thyroid dysfunction among pregnant women and its effect on maternal and fetal outcome. *J Obstet Gynecol India*. 2014; 64(2): 105- 10.
15. Mannisto T, Vaarasmäki M, Pouta A, et al. Thyroid dysfunction and autoantibodies during pregnancy as predictive factors of pregnancy complications and maternal morbidity in later life. *J Clin Endocrinol Metab*. 2010; 95(3): 1084- 94.
16. Männistö T, Vääräsmäki M, Pouta A, et al. Perinatal outcome of children born to mothers with thyroid dysfunction or antibodies: a prospective population-based cohort study. *J Clin Endocrinol Metab*. 2009; 94(3): 772- 79.

17. Turunen S, Vääräsmäki M, Männistö T, et al. Pregnancy and perinatal outcome among hypothyroid mothers: a population-based cohort study. *Thyroid*. 2019; 29(1): 135- 41.
18. Naderi T, Honarvar Z, Bahrampor A, Yosefzadeh G. The prevalence of hypothyroidism based on risk factors in pregnant women referred to Shahid Dadbin Clinic, Kerman, Iran. *J Kerman Univ Med Sci*. 2012; 19(3): 225-32. [In Persian]
19. Klein RZ, Haddow JE, Falx JD, et al. Prevalence of thyroid deficiency in pregnant women. *Clin Endocrinol*. 1991; 35(1): 41- 6.
20. Cleary-Goldman J, Malone FD, Lambert-Messerlian G, et al. Maternal thyroid hypofunction and pregnancy outcome. *Obstet Gynecol*. 2008; 112(1): 85-92.
21. Cakmak BD, Turker UA, Temur M, Ustunyurt E. Pregnancy outcomes of antibody negative and untreated subclinical hypothyroidism. *J Obstet Gynaecol Res*. 2019; 45(4): 810- 16.
22. Shinohara DR, Santos Th, de Carvalho HC, et al. Pregnancy complications associated with maternal hypothyroidism: a systematic review. *Obstet Gynecol Survey*. 2018; 73(4): 219- 30.
23. Chen L-M, Du W-J, Dai J, et al. Effects of subclinical hypothyroidism on maternal and perinatal outcomes during pregnancy: a single-center cohort study of a Chinese population. *PLoS One*. 2014; 9(10): e109364.
24. Stohl HE, Ouzounian J, Rick A-M, Hueppchen NA, Bienstock JL. Thyroid disease and gestational diabetes mellitus (GDM): is there a connection? *J Matern Fetal Neonatal Med*. 2013; 26(11): 1139- 42.
25. Lee SY, Cabral HJ, Aschengrau A, Pearce EN. Associations Between Maternal Thyroid Function in Pregnancy and Obstetric and Perinatal Outcomes. *J Clin Endocrinol Metab*. 2020; 105(5): e2015- e23.
26. Luisi S, Giorgi M, Riggi S, Messina G, Severi FM. Neonatal outcome in pregnancy hypotiroidee women. *Gynecol Endocrinol*. 2020; 36(9): 772-75