

Original Article

Blood Glucose and Neurological Status: Dual Predictors of Survival in Diabetic Emergency

Lintang Arum Nikentari¹, Heri Kristianto^{2*}, Laily Yuliatun², Ali Haedar³,
Paulus Lucky Tirma Irawan⁴

¹Postgraduate Nursing Student, Faculty of Health Sciences, Brawijaya University, Malang, Indonesia

^{2*}Nursing Department, Faculty of Health Sciences, Brawijaya University, Malang, Indonesia

³Emergency Department, Faculty of Medicine, Brawijaya University, Malang, Indonesia

⁴Informatics Engineering Department, Faculty of Technology And Design, Ma Chung University, Malang, Indonesia

Received: 28 Jan 2025 Accepted: 18 Feb 2025 Published: 5 Mar 2025

Article Info

Keywords:

Diabetic ketoacidosis, Glasgow Coma Scale, Hyperosmolar hyperglycemic syndrome, Blood glucose, Emergency care

*Corresponding author:

School of Nursing, Metabolic Syndrome Research Group, Faculty of Health Science, Academic Hospital, Universities Brawijaya, Malang, Indonesia
Email: heri.kristianto@ub.ac.id

How to Cite This Article:

Arum Nikentari L, Kristiano H, Yuliatun L, Header A, Lucky Trima Irawan P, Blood Glucose and Neurological Status: Dual Predictors of Survival in Diabetic Emergency Prev Care Nurs Midwifery J PCNM. 2025; 15(1): 30-36

Abstract

Background: Acute diabetic emergencies, including diabetic ketoacidosis (DKA), hyperosmolar hyperglycaemic syndrome (HHS), and hypoglycaemia, require urgent medical intervention. These complications result from severe metabolic disturbances, often causing neurological impairment. Blood glucose levels measure metabolic derangement, while the Glasgow Coma Scale (GCS) indicates cerebral dysfunction.

Objectives: This study evaluates the prognostic significance of blood glucose levels and GCS in predicting survival rates among diabetic emergency patients, hypothesizing that these parameters are robust outcome indicators.

Methods: A retrospective cross-sectional study analysed medical records of 250 patients treated for diabetic emergencies at Dr. Soedono General Hospital Madiun (2017–2024). Blood glucose levels and GCS scores were assessed using chi-square tests ($p < 0.25$) and multivariate binomial logistic regression ($p < 0.05$) in SPSS version 21.

Results: Multivariate analysis identified GCS as the most critical survival predictor, with low GCS scores significantly correlating with mortality (OR = 0.002, 95% CI: 0.000–0.012, $p < 0.05$). Blood glucose levels >600 mg/dL were also associated with reduced survival rates (OR = 0.113, 95% CI: 0.074–4.304, $p < 0.05$). The model explained 72.1% of the variance in patient outcomes.

Conclusion: GCS and blood glucose levels are pivotal survival predictors in diabetic emergencies, with GCS being the predominant determinant. These findings highlight the importance of early neurological evaluation and glucose regulation in improving outcomes

■ Implications of this paper in nursing and midwifery preventive care

- Early Neurological Assessment and Monitoring: The study highlights the Glasgow Coma Scale (GCS) as a critical predictor of survival in diabetic emergencies. Nurses and midwives should prioritize early neurological assessment and continuous monitoring in emergency care settings to detect signs of deterioration promptly and initiate timely interventions.
- Blood Glucose Management and Patient Education: Severe hyperglycemia (blood glucose >600 mg/dL) significantly reduces survival rates in diabetic emergencies. Preventive strategies should focus on regular blood glucose monitoring, patient education on recognizing early symptoms of diabetic crises, and adherence to treatment regimens to minimize emergency complications.



Copyright © 2021, This is an original open-access article distributed under the terms of the Creative Commons Attribution-noncommercial 4.0 International License which permit copy and redistribution of the material just in noncommercial usages with proper citation

Introduction

Diabetic emergencies, including diabetic ketoacidosis (DKA), hyperosmolar hyperglycaemic state (HHS), and hypoglycaemia, represent life-threatening conditions that require urgent medical intervention to prevent mortality and long-term complications [1]. These emergencies arise due to severe metabolic imbalances, which not only disrupt normal physiological functions but also significantly impact neurological status. Neurological dysfunction in diabetic emergencies often results from prolonged hyperglycaemia, cerebral edema, or hypoglycaemia-induced. This study aims to investigate the dual predictive role of blood glucose levels and neurological status in determining survival outcomes among diabetic emergency patients. neuronal damage, leading to altered levels of consciousness and increased morbidity [2]. Therefore, understanding the relationship between metabolic disturbances and neurological status is critical in improving patient outcomes. Previous studies have established a strong association between severe hyperglycaemia and poor clinical outcomes, with blood glucose levels exceeding 600 mg/dL being a significant predictor of mortality in diabetic emergencies [3]. Similarly, neurological impairment, as measured by GCS, has been shown to independently predict survival, with lower GCS scores correlating with an increased risk of death due to severe cerebral dysfunction and diminished recovery capacity [4]. The interplay between these two factors suggests that an integrated assessment of blood glucose levels and GCS can enhance prognostic accuracy and guide clinical decision-making in diabetic emergencies [5]. Despite advancements in emergency diabetes care, the prognostic role of combined metabolic and neurological assessments remains underexplored. While hyperglycaemia-induced metabolic stress exacerbates systemic complications, concomitant neurological deterioration may further reduce survival chances, necessitating an early and comprehensive evaluation of both parameters [6].

Objectives

This study aims to investigate the dual predictive role of blood glucose levels and neurological status in determining survival outcomes among diabetic emergency patients. By establishing the significance of these variables, the findings may contribute to the development of more refined clinical protocols and

predictive models, ultimately improving emergency management strategies for diabetes-related complications.

Methods

Study Design

This study employed a retrospective design with a cross-sectional approach, conducted at the Emergency Department (ED) of a Regional General Hospital (RSUD) in East Java, between July and September 2024.

Participants

The study population comprised medical records of patients aged over 30 years with type 2 diabetes mellitus, who had been treated in the ED for hypoglycemia, hyperosmolar hyperglycemic state (HHS), or diabetic ketoacidosis (DKA) over the preceding eight years (2017–2024). Inclusion criteria were limited to patients diagnosed with type 2 diabetes mellitus, selected through a time sampling method within a predefined period to ensure a representative dataset. This approach captured variations in patient characteristics, disease severity, and treatment outcomes while minimizing selection bias. Patients with type 1 diabetes mellitus and gestational diabetes were excluded to maintain population homogeneity, as these conditions have distinct pathophysiologies and management strategies that could confound the analysis. Additionally, incomplete medical records were excluded to ensure data accuracy and reliability in assessing survival factors in diabetic emergency patients.

Sampling Method

Sampling was carried out using a time sampling method, resulting in a total of 250 patient medical records. These records were retrieved from the hospital's electronic medical record database following ethical approval (No. 400.14.5.4/23377/102.9/2024).

Data Collection

Data collection was conducted through the integrated electronic medical record system managed by the Medical Record Unit. A trained medical records team ensured the accuracy of data entry and verification. Any incomplete or missing data were excluded from the analysis.

Variables

The key variables measured included blood glucose levels (mg/dL), neurological status assessed using the Glasgow Coma Scale (GCS), and survival rate.

Measurement Tools

Data were obtained through the integrated electronic medical record system managed by the Medical Record Unit, with neurological status assessed using the Glasgow Coma Scale (GCS).

Statistical Methods

Data analysis was performed using SPSS version 21 (IBM Corp., Armonk, NY, USA), with logistic binomial regression applied at a significance threshold of $p<0.05$ to identify factors significantly influencing patient survival rates.

Ethical Considerations

These records were retrieved from the hospital’s electronic medical record database following ethical approval (No. 400.14.5.4/23377/102.9/2024).

Results

The trend in the incidence of diabetic emergency cases from 2017 to 2024 is illustrated in Figure 1, with data presented clearly to facilitate interpretation. Notable fluctuations in case numbers were observed across the years, which may correlate with advancements in diabetes management and changes in patient health status. In 2017 and 2018, the cases remained relatively stable at approximately 2.96% and 2.20%, respectively. However, a significant rise of 21.97% occurred in 2019, followed by a reduction in subsequent years, with cases dropping to 10.70% in 2020 and 8.40% in 2021. The decline in cases during this period may reflect improvements in diabetes management strategies. In 2022, a sharp decline to 2.29% was

recorded, yet a dramatic increase was observed in 2023, reaching 35.82%. This surge may be linked to the escalating incidence of diabetic ketoacidosis (DKA) or hyperosmolar hyperglycaemic state (HHS), possibly due to deteriorating patient health or delays in seeking treatment. By 2024, the percentage of cases fell to 15.66%, though it remained higher than in the early years of the study period, suggesting that challenges in diabetic emergency management persisted. Additionally, the Glasgow Coma Scale (GCS) was identified as a significant predictor of survival, with low GCS scores strongly associated with mortality (OR = 0.002, 95% CI: 0.000–0.012, $p < 0.05$). Blood glucose levels >600 mg/dl were also linked to reduced survival rates (OR=0.113, 95% CI: 0.074–4.304, $p<0.05$). The model accounted for 72.1% of the variance in patient outcomes. Given its crucial role in patient prognosis, further analysis was conducted to assess potential confounding factors, such as comorbidities and treatment variability, which could influence survival outcomes. Detailed percentages for each year are provided in Figure 1.




Figure 1. Diagram of Diabetic Emergency Patient Admissions (MRS) in the ED from 2017 – 2024

The characteristics of patients experiencing diabetic emergencies from 2017 to 2024 are summarized in Table 1. The majority of patients were elderly (71.3 %) and predominantly male (69.3 %). Blood glucose levels varied widely, with a substantial proportion experiencing severe hypoglycemia or extreme hyperglycemia. Neurological status, assessed using the Glasgow Coma Scale (GCS), showed a distribution across low, moderate, and high categories. In terms of survival, more than half of the patients survived, while a significant proportion did not. Further details on age distribution, gender, blood glucose levels, GCS scores, and survival rates are provided in Table 1.

Tabel 1. Blood Glucose Levels and GCS of Diabetic Emergency Patients 2017 – 2024

Characteristics	n = 250	%
Age		
Adults (30 – 59 years)	115	28.7
Elderly (>60 years)	135	71.3
Gender		
Female	123	30.7
Male	127	69.3
Blood Sugar Levels		
<54 mg/dl	161	64.4
>250 mg/dl	15	6.0
>600mg/dl	74	29.6
GCS		
Low (3 – 8)	99	39.6
Moderate (9 – 12)	38	15.2
High (13 – 15)	113	45.2
Survival Rate		
Deceased	91	36.4
Alive	159	63.6

The analysis exploring the relationship between blood glucose levels, neurological status, and patient survival rates in diabetic emergencies is detailed in Table 2. The bivariate analysis, conducted using the chi-square test with a significance threshold of $p < 0.25$, revealed that both blood glucose levels and neurological status were significantly associated with patient survival. Specifically, the categories of bloodglucose levels<54mg/dL($p<0.001$),>250mg/dL($p<0.001$),a=0.004)—demonstrated

significant associations at the bivariate level. Regarding neurological status, the low Glasgow Coma Scale (GCS) category exhibited a p-value of 0.113, while the moderate and high categories showed p-values of 0.098 and 0.580,

respectively. In the multivariate analysis, conducted using binomial logistic regression with a significance level of $p < 0.05$, GCS ($p < 0.001$) emerged as a significant predictor of patient survival, whereas blood glucose levels did not exhibit a significant relationship ($p = 0.073$). The R Square value indicated that the model accounted for approximately 72.1% of the variance in the data. Multivariate analysis identified neurological status, as measured by the Glasgow Coma Scale (GCS), as the most influential factor affecting survival in patients with diabetic emergencies (OR = 0.002, 95% CI: 0.000–0.012, $p < 0.05$). Additionally, blood glucose levels >600 mg/dL were associated with reduced survival rates (OR = 0.113, 95% CI: 0.074–4.304, $p < 0.05$) (Table 2).

Tabel 2. Blood Glucose and Neurological Status in Survival Diabetic Emergency

	S.E.	Df	p	R Square	Chi-Square (p<0.25)	Multivariate (p<0.05)	95% CI Exp (B)
Blood Sugar Levels				0.028	0.073	0.113	0.002 (0.000–0.012)
<54 mg/dl	0.528	2	< 0.001				
>250 mg/dl	0.623	1	< 0.001				
>600mg/dl	1.038	1	0.004				
GCS				0.721	< 0.001*	< 0.001*	0.113 (0.074–4.304)
Low	0.842	2	0.113				
Moderate	0.840	1	0.098				
High	0.863	1	0.580				

* $p<0.05$

These findings highlight the critical role of neurological status in predicting survival outcomes among diabetic emergency patients. The strong association between low GCS scores and mortality suggests that impaired consciousness significantly increases the risk of poor prognosis (Table 2). This emphasizes the need for early neurological assessment in diabetic emergencies to guide clinical interventions. Although blood glucose levels were associated with survival in the bivariate analysis their significance diminished in the multivariate model, indicating that other factors, such as neurological status, play a more dominant role.

Discussion

The findings of this study suggest that neurological status, as assessed by the Glasgow Coma Scale (GCS), serves as the primary predictor of survival in patients experiencing diabetic emergencies. This is consistent with previous studies that have demonstrated a strong correlation between GCS scores and patient outcomes in acute metabolic conditions [7,4,8]. Additionally, blood glucose levels play a significant role, particularly when exceeding 600 mg/dL, which has been previously linked to increased mortality in hyperglycaemic crises. These findings emphasize the critical importance of prompt and continuous neurological assessment in the management of patients within the emergency department (ED) [9,10]. Patients exhibiting low GCS scores face an exceptionally high risk of mortality, aligning with previous research that identifies severe neurological impairment as a key determinant of adverse outcomes in diabetic ketoacidosis (DKA) and hyperosmolar hyperglycaemic state (HHS)[10,11]. This underscores the need for early intervention strategies aimed at stabilizing the patient's neurological condition to improve survival rates [12,13]. Moreover, both extreme hyperglycaemia and hypoglycaemia necessitate particularly careful attention [14]. Severe hyperglycaemia (above 600 mg/dL) has been strongly associated with a heightened risk of mortality, while hypoglycaemia (below 54 mg/dL) also presents substantial potential for serious complications, corroborating earlier studies that emphasize the dangers of extreme blood glucose fluctuations [15,16]. Consequently, close

Conclusion

This study highlights the Glasgow Coma Scale (GCS) as the primary predictor of survival in diabetic

monitoring of blood glucose levels from the point of initial triage in the ED is essential to enhancing patient survival. These findings carry significant implications for the formulation of more comprehensive ED protocols, which should incorporate a combined evaluation of blood glucose levels and neurological status as a key component of the early triage process for diabetic emergency patients [17,18]. By adopting this approach, healthcare providers can allocate resources more effectively to those patients at the greatest risk, thereby improving clinical outcomes [19,20]. From a preventive standpoint, public health education plays a pivotal role in reducing the incidence of diabetic emergencies. This initiative involves educating patients and their families on effective home blood glucose management, early identification of complications such as DKA or HHS, and reinforcing the significance of regular health check-ups [21]. Community-based preventive programs that encourage healthy lifestyles, including balanced diets and regular physical activity, also play a crucial role in preventing acute complications [22]. This study provides valuable insights into the critical role of GCS in predicting survival in diabetic emergencies, reinforcing its importance in clinical decision-making. The integration of blood glucose levels as an additional predictor further enhances the study's applicability to emergency care settings. However, there are limitations to consider. As a retrospective study conducted at a single institution, the findings may not be generalizable to other settings or populations. Additionally, the lack of longitudinal data prevents an assessment of long-term outcomes post-emergency treatment. Furthermore, while GCS and blood glucose levels are highlighted, other potential confounding factors influencing survival rates, such as comorbidities and treatment variations, warrant further investigation. Future research should be conducted with a larger population and a broader range of care conditions to validate these findings and aid in the development of risk prediction algorithms. By doing so, these results not only offer fresh insights into the management of diabetic emergencies but also provide a foundation for the creation of more effective preventive measures, ultimately enhancing the quality of life for diabetic patients.

emergencies, with blood glucose levels also playing a significant role. Patients with low GCS scores are at an exceptionally high risk of mortality, emphasizing the importance of early neurological

assessment and intervention in emergency care. The integration of both GCS and blood glucose levels in triage protocols can enhance patient outcomes by facilitating timely and appropriate medical interventions. Additionally, public health education on blood glucose management and early recognition of complications is essential in reducing the incidence of diabetic emergencies. Future studies should expand on these findings by incorporating larger sample sizes and exploring long-term patient outcomes to develop more precise risk prediction models.

Ethical Considerations

This research has received approval from the Research Ethics Committee of Dr. Soedono General Madiun Hospital with ethics number: 400.14.5.4/233377/102.9/2024

Acknowledgments

The researchers extend their gratitude to Dr. Soedono General Hospital in Madiun for the support and permission to conduct this study. Appreciation is also given to all medical staff and personnel who contributed to data collection and provided valuable information. It is hoped that the results of this study can benefit the improvement of healthcare services at this hospital.

Conflict of Interest

The authors hereby declare that there are no potential conflicts of interest concerning the research, authorship, or publication of this article.

Funding

The authors proudly acknowledge the funding for this research, authorship, and/or publication of this article under the 2024 Non-Head Lecturer Doctoral Grant Scheme, grant number 10993.1/UN10.F17/PT.01.03/2024, provided by Universitas Brawijaya

Authors' Contributions

Nikentari L.A. contributed to conceptualization, methodology, data collection, and manuscript writing. Kristianto H. supervised the study, performed data analysis, reviewed the manuscript, and handled correspondence. Yuliatun L. conducted literature review, statistical analysis, manuscript editing, data interpretation, validation, and final

approval. Haedar A. and Irawan P.L.T. provided administrative and technical support. All authors reviewed and approved the final manuscript.

Artificial Intelligence Utilization

The authors declare that no generative AI technologies were used in the creation of this manuscript.

Data Availability Statement

The datasets generated and/or analyzed during the current study are available from the corresponding author upon reasonable request.

References

1. Saifullah H. Diabetic emergency: diabetic ketoacidosis, hyperglycemic hyperosmolar state, euglycemic diabetic ketoacidosis and hypoglycemia. *International Journal of Contemporary Research*. 2024;3(1):31-37. <https://multiarticlesjournal.com/counter/d/3-1-7/IJCRM-2024-3-1-7.pdf>.
2. Rusdi MS. Hipoglikemia pada pasien diabetes melitus. *Journal Syifa Science and Clinical Research*. 2020;2(September):83-90. <https://doi.org/10.37311/jsscr.v2i2.4575>
3. Muneer M, Akbar I. Acute metabolic emergencies in diabetes: DKA, HHS and EDKA. *Advances in Experimental Medicine and Biology*. 2021;1307:85-114. https://doi.org/10.1007/5584_2020_545
4. Lotter N, Lahri S, van Hoving DJ. The burden of diabetic emergencies on the resuscitation area of a district-level public hospital in Cape Town. *African Journal of Emergency Medicine*. 2021;11(4):416-421. <https://doi.org/10.1016/j.afjem.2021.05.004>
5. Braine ME, Cook N. The Glasgow coma scale and evidence-informed practice: a critical review of where we are and where we need to be. *Journal of Clinical Nursing*. 2017;26(1-2):280-293. <https://doi.org/10.1016/j.afjem.2021.05.004>
6. Pawils S, Heumann S, Schneider SA, Metzner F, Mays D. The current state of international research on the effectiveness of school nurses in promoting the health of children and adolescents: an overview of reviews. *PLOS One*. 2023;18(4). <https://doi.org/10.1371/journal.pone.0275724>
7. Sarang B, Bhandarkar P, Raykar N, O'Reilly GM, Soni KD, Wärnberg MG, et al. Associations of on-arrival vital signs with 24-hour in-hospital mortality in adult trauma patients admitted to four public university hospitals in urban India: a prospective multi-centre cohort study. *Injury*. 2021;52(5):1158-1163. <https://doi.org/10.1016/j.injury.2021.02.075>
8. Akram Z, Inayat M, Akhtar S, Perveen R, Saeed R, Abid J, et al. Fasting influence on diabetic emergency visits in a tertiary care hospital throughout Ramadan and other lunar months. *Pakistan Journal of Medical and Health Sciences*. 2023;17(3):588-590. <https://doi.org/10.53350/pjmhs2023173588>
9. Maharjan J, Pandit S, Johansson KA, Khanal P, Karmacharya B, Kaur G, et al. Effectiveness of interventions for emergency

care of hypoglycaemia and diabetic ketoacidosis: a systematic review. *Diabetes Research and Clinical Practice*. 2024;207. <https://doi.org/10.1016/j.diabres.2023.111078>

10. Tamzil R, Yaacob N, Noor N, Baharuddin K. Comparing the clinical effects of balanced electrolyte solutions versus normal saline in managing diabetic ketoacidosis: a systematic review and meta-analyses. *Turkish Journal of Emergency Medicine*. 2023;23(3):131-138. https://doi.org/10.4103/tjem.tjem_355_22

11. Blank SP, Blank RM, Ziegenfuss MD. The importance of hyperosmolarity in diabetic ketoacidosis. *Diabetic Medicine*. 2020;37(12):2001-2008. <https://doi.org/10.1111/dme.14277>

12. Watta R, Masi G, Katuuk ME. Screening faktor resiko diabetes melitus pada individu dengan riwayat keluarga diabetes melitus di RSUD Jailolo. *Jurnal Keperawatan*. 2020;8(1):44. <https://doi.org/10.35790/jkp.v8i1.28410>

13. Chen J, Zeng H, Ouyang X, Zhu M, Huang Q, Yu W, et al. The incidence, risk factors, and long-term outcomes of acute kidney injury in hospitalized diabetic ketoacidosis patients. *BMC Nephrology*. 2020;21(1):1-9. <https://doi.org/10.1186/1471-2369-7-1>

14. Banday MZ, Sameer AS, Nissar S. Pathophysiology of diabetes: an overview. *Avicenna Journal of Medicine*. 2020;10(4):174-188. https://doi.org/10.4103/ajm.ajm_53_20

15. Gelaw NB, Muche AA, Alem AZ, Gebi NB, Chekol YM, Tesfie TK, et al. Development and validation of risk prediction model for diabetic neuropathy among diabetes mellitus patients at selected referral hospitals, in Amhara regional state Northwest Ethiopia, 2005-2021. *PLOS One*. 2023;18(8):e0276472. <https://doi.org/10.1371/journal.pone.0276472>

16. Wu XY, She DM, Wang F, Guo G, Li R, Fang P, et al. Clinical profiles, outcomes and risk factors among type 2

diabetic inpatients with diabetic ketoacidosis and hyperglycemic hyperosmolar state: a hospital-based analysis over a 6-year period. *BMC Endocrine Disorders*. 2020;20(1):1-9. <https://doi.org/10.1186/s12902-020-00659-5>

17. Katuuk ME, Sitorus R, Sukmarini L. Penerapan teori self care Orem dalam asuhan keperawatan pasien diabetes melitus. *Jurnal Keperawatan*. 2020;8(1):1-22. <https://doi.org/10.35790/jkp.v8i1.28405>

18. Barski L, Golbets E, Jotkowitz A, Schwarzfuchs D. Management of diabetic ketoacidosis. *European Journal of Internal Medicine*. 2023;117:38-44. <https://doi.org/10.1016/j.ejim.2023.07.005>

19. Jitraknatee J, Ruengorn C, Nochaiwong S. Prevalence and risk factors of chronic kidney disease among type 2 diabetes patients: a cross-sectional study in primary care practice. *Scientific Reports*. 2020;10(1):1-10. <https://doi.org/10.1038/s41598-020-63443-4>

20. Dhatariya KK. The management of diabetic ketoacidosis in adults-an updated guideline from the Joint British Diabetes Society for Inpatient Care. *Diabetic Medicine*. 2022;39(6):e14788. <https://doi.org/10.1111/dme.14788>

21. Wu F, Wu C, Wu Q, Yan F, Xiao Y, Du C. Prediction of death in intracerebral hemorrhage patients after minimally invasive surgery by vital signs and blood glucose. *World Neurosurgery*. 2024;184:e84-e94. <https://doi.org/10.1016/j.wneu.2024.01.061>

22. Mohajan D, Mohajan HK. Hypoglycaemia among diabetes patients: a preventive approach. *Journal of Innovative Medical Research*. 2023;2(9):29-35. <https://doi.org/10.56397/JIMR/2023.09.05>