Preventive Care in Nursing and Midwifery Journal 2018; 8(3): 35-41

A Comparative Study of the Life Quality and Style of the Healthy and Asthma Suffering People Referring to Zanjan Province Hospitals

Rajabi S¹, <u>Aghvamy MS</u>^{1*}, Bahraminejad N², Fallah R³

^{*1}Department of Nursing, School of Nursing and Midwifery, Zanjan University of Medical Sciences, Zanjan, Iran

²Department of Nursing, Social Determinant of Health Research Center, Assistant Professor, School of Nursing and Midwifery, Zanjan University of Medical Science, Zanjan, Iran

³Department of Biostatistics and Epidemiology, Assistant Professor, School of Medicine, Zanjan University of Medical Sciences, Zanjan, Iran

*Corresponding Author: Department of Nursing, School of Nursing and Midwifery, Zanjan University of Medical Sciences, Zanjan, Iran

> *Email:* aghvamym@zums.ac.ir Received: 31 Aug 2019 Accepted: 24 Dec 2019

Abstract

Background: Today, asthma is one of the most prevalent chronic respiratory diseases, involving a large number of people and communities. Research show that the disease overshadows the social, psychological and even economic life and performance of the patients.

Objectives: This study was conducted in Zanjan Province hospitals with the purpose of comparing the life style and quality of life asthma patients with healthy people.

Methods: This study is descriptive-comparative in nature, focusing on two groups of individuals (healthy people and those suffering from asthma) using convenience sampling method to serve the goals of the study. The instruments used for collection of data were the World Health Organization Quality of Life (WHOQOL) questionnaire and the Life Style Questionnaire (LSQ). The data were analyzed using the SPSS software version 16.

Results: The demographic data relating to both groups (n=50 for each group) did not show any difference of statistically significant. The mean scores of the life style of the healthy people and the asthma patients were 149.34 and 136.48, respectively (p<0.05). Furthermore, the mean scores of the Quality of Life (QOL) of healthy people and asthmatics were 36.260 and 29.205, respectively (p<0.05).

Conclusion: The life style and QOL of asthma patients are deteriorated to a large extent. Therefore, it is recommended that educational interventions to improve the life style and QOL of patients included in their care plan.

Keywords: asthma, quality of life, lifestyle, asthma patients

Introduction

Asthma is a disease characterized by inflammation of the airways, lungs distinguishably the narrowing of the airways and blocking of the airflow and leading to symptoms of suffocation, hard breathing and change in indices of the patient's respiratory test [1]. Asthma is one of the most prevalent chronic diseases around the world [2], leaving drastic impact on the lives and performance of patients influencing collective activities while and reversing even the course of social, psychological

and economic performance of patients. This will afterwards cause other intricacies and problems in the patient [3]. As many as 300 million people in the world seem to have been suffering from the disease [4]. As far as dominance of the disease in Iran is concerned, it is five percent to 15 percent on the average [5]. Besides causing high rate of mortality, the disease affects patients' communication, and different aspects of their life like family dimensions and social activities, while restricting patients' physical activity [6-8]. For the time being no ultimate medication is supposed for asthma; however, appropriate medicinal treatment, better lifestyle, and healthrelated habits are believed to be the best and the most influential mechanisms proposed to check progress of the disease [9,10]; this way any improvement in the Quality of Life (QOL) criteria will subsequently yield lesser amount of the symptoms of asthma [11]. By 2020, 40 to 50 percent of the lifestyle related non-contagious diseases would account for seven out of 10 death cases in the developing countries [12,13]. Statistics on global rate of mortality, caused by chronic respiratory diseases such as asthma show that asthma stands as the fourth major cause of deaths worldwide [14]. The more attention paid to lifestyle and its improvement would result in avoidance of other health disorders like depression [15]; diabetes [16,17]; obesity; problems with absorption of vitamin D; and heart disease as a result of asthma [18,19].

Depending on the cultural conditions and beliefs of people in each geographic region, diseases can have different effects on patients' QOL and lifestyle. On this basis and because no research has been done on the QOL and lifestyle of the asthma patients in Zanjan Province, this study aims to compare the life style and QOL of asthma patients with healthy people.

Methods

This research is descriptive comparative in nature, carried out on the asthma patients referring to selected hospitals of Zanjan Province from 2017 to 2018. The study was done after receiving the code of ethics from the Ethics Committee of Zanjan University of Medical Sciences (irzums.rec.1397.90) and once the participants signed the form of informed consent. The participants were selected on the convenience sampling method. The healthy people was selected from the patients' companions referred to the lung ward. Since the life style samples outnumbered the OOL samples, the level of significance was decided to be five percent and the test power was set to be 90 percent with respect to the sample size.

The research setting were internal wards of Valiasr Hospital in Zanjan, Army Hospital (554) of Zanjan, Zanjan Social Security Hospital, Al Ghadir Hospital (Abhar) Social Security Hospital (Abhar), Boali Ali Sina Hospital of Khorramdareh, Razi Hospital of Mahneshan, Amiralmomenin Hospital of Khodabandeh, Shohadaye Tarom Hospital and Isar Ijrud Hospital.

Characteristics of study participants included age range of 40-60 years, confirmation of disease by a physician, lack of psychological problems and substance abuse. The method of convenience sampling continued until having required samples. Meanwhile, the instruments for collection of data included: A questionnaire to obtain demographic information on the respondents' sex, age, educational status, job, income, marital status, medication follow-up status, Smith Miller Lifestyle Questionnaire (LSQ) and an abbreviated 26-item version of the World Health Organization QOL questionnaire (WHOQOL-BREF).

The WHOQOL questionnaire , designed by the World Health Organization (WHO) in 1998 to assess the QOL, consists of 26 questions, including 24 questions to assess the four domains of physical health (7 questions), mental health (6 questions), social relations (3 questions) and environmental health (8 questions) and 2 questions to assess the overall QOL domain. Nejat et al. (2006) validated the scale in Iran context and obtained the alpha coefficient of the questionnaire for healthy population. The alpha coefficient for the physical health domain was 0.70, for mental health was 0.73, for social relations was 0.55 and for environmental communication was 0.84 [20].

The LSQ has a total of 70 questions. The questionnaire consists of 10 factors or components including physical health, exercise and health, weight and nutrition control, disease prevention, psychological health, spiritual health, social health, drug consumption avoidance, narcotic drugs, alcohol, accident prevention and environmental health. This questionnaire is scored on a Likert scale as "Always" (3), "Usually" (2), "Sometimes" (1), and "Never" (0). High scores in each of the components and the total questionnaire indicate a more appropriate lifestyle [21]. The reliability and validity of the questionnaire had previously been estimated by Lali et al. [21]. The SPSS software version 16 was used to analyze the data gained from the questionnaire filled up by the respondents. The means of the QOL and lifestyle variables

pertinent to the two groups under study were compared using the descriptive statistics and ttest.

Results

There were a total of 100 participants, including 61 men and 39 women, in this study. 17% of them were single, 76% were married, 7% were divorced. Out of the population, 62% were urban dwellers and 38% were rural citizens. Economically speaking, 41 percent of the individuals were in poor economic status, 42 percent were in the middle class, and 17 percent had good economic status. As far as their age is

concerned, 26 of the individuals were below 40, 42 fell within the age range of 40 to 60; 32 of the individuals were above 60. As for their educational status, 22 percent of them were illiterate, 36 percent of them were under a diploma, 26 percent held diploma and 36% were college educated. From standpoint of their employment, 11 percent were state employees, 22 percent were freelancers, 16 percent were laborers, 22 percent were housewives and 29 percent were unemployed. Table 1 shows information on the demographic characteristics of the members of study groups.

Table 1: Frequency Distribution of Participants by Group Membership and Demographic Factors

Demographic Variables		Asthma Patients N=50	Healthy People N=50	
Condon	Male	29	32	
Gender	Female	21	18	
	Single	11	6	
Marital	Married	36	40	
Status	Divorced	3	1	
	Widow	-	3	
Address	Urban	30	32	
Address	Rural	20	18	
The	Low Class	16	25	
Economic	Middle Class	25	17	
Status	Top Class	9	8	
.	<40	20	6	
Age (In Vears)	40 to 59	17	25	
Tears)	>=60	13	19	
	Illiterate	8	14	
Educational	High School	10	16	
Status	Diploma	13	13	
	Bachelor and Above	19	7	
	Employee	8	3	
	Free	11	11	
Job Status	Manual Worker	9	7	
	Housewives	11	11	
	Unemployed	11	18	

According to Table 2, independent t-test showed a significant difference between the mean scores of the asthmatic-non-asthmatic patients' lifestyle in

terms of physical health, exercise and health, psychological health, social health and environmental health (P<0.05).

			Group			
Variable		With Asthma		Healthy		P Value
		Average	Standard Deviation	Mean	Standard Deviation	(T-Test)
Dimen - sions - of Life - Style -	Physical Health	16.02	2.27	18.28	2.50	0.005
	Exercise and Health	13.34	3.89	15.80	3.24	0.005
	Weight Control and Nutrition	12.32	2.89	12.60	2.74	0.621
	Disease Prevention	12.28	2.71	13.14	2.89	0.129
	Psychological Health	14.54	2.65	16.58	2.72	0.005
	Mental Health	13.66	2.20	13.92	2.16	0.554
	Social Health	13.42	2.75	16.28	1.75	0.005
	Avoid Medicines	13.38	2.52	12.72	2.37	0.181
	Accident Prevention	15.18	2.53	15.88	1.99	0.128
	Environmental Health	12.34	2.38	14.14	2.32	0.005
	Total	136.48		149.34		

Table 2: Independent t-test results for lifestyle variable and its dimensions by study group

Table 3 shows that general mean and all dimensions of QOL were significantly lower in

asthmatic patients than non-asthmatic patients (P<0.05) based on the independent t-test results.

Table 3: Independent T-Test Results for Quality of Life Variable and It's Dimensions by Study Group

		Group				P Value
Variable		With Asthma		Healthy		
		Average	Standard Deviation	Mean	Standard Deviation	(T-Test)
Dimensions of Quality of Life	Physical Health	45/34	21/86	65/42	15/01	0/005
	Mental Health	48/41	14/54	59/04	14/83	0/005
	Community Relations	63/01	18/73	77/72	15/57	0/005
	Environmental Health	48/53	19/51	58/18	18/20	0/012
	Total	205.29		260.36		

Discussion

The results of this study showed difference between the mean scores of most aspects of lifestyle and QOL in the asthmatic and nonasthmatic patients. A longitudinal study showed that lowering amount of physical activities and a low volume of daily physical activities, as well as a selection of sedentary activities such as watching television and computer games were associated with worsening clinical manifestations of asthma [22]. The results of a study in Taif region of Saudi Arabia showed that the quality of life of children and adults with controlled asthma was higher than those of the patients with uncontrolled asthma [23]. Amira et al. (2017) showed that levels of asthma control in individuals (activities, symptoms, and environment) had significant relationship with the

quality of life of the individuals suffering asthma [11].

These results can be interpreted as asthma causing limitations in one's social activities to the extent that the individuals will have less social presence and activities than others [24]. The asthma patients are proved to have been experiencing emotional disorders, anxiety, hostility, and other psychological factors, which influence their happiness [25]. Many asthmatic patients are also shown to have low level of concentration because of depression, as the low concentration and depression jointly downgrade their quality of life [26]. The quality of life lowered on mediation of the associated mental disorders prompts poor function of the patients' lungs. The low quality of life in turn provokes ever growing severity of illness, frequent hospitalizations, burnout, fatigue,

and readiness to behavioral problems and restricted activities [26].

The condition of quality of life illustrated that the mean score of asthmatic patients was lower than the healthy individuals. Salman Yazdi et al. reported the mean score of the QOL of girls standing at 50.56 and that of boys with asthma at 36.02 [27]. Arash et al. also proved that asthmatic patients in majority of the wards had a relatively favorable QOL (58.3%) [28], the difference between results of this study and those of other studies might be cultural and age differences between the two groups.

Results of the analysis of showed that the lifestyle scores of the healthy individuals were higher and better than the asthmatic patients. Rouhafza et al. studied the Chronic Obstructive Pulmonary Disease (COPD) patients and proved that the lifestyle scores of the COPD patients signed their unhealthy life style [29]. Hassanpour et al. showed that there was significance difference in the scores of the minor and medium phases of the mental disorder domain scores of qualify of life in COPD patient. As for the symptoms' domain, there was a significant difference between the total score and the score of the two severe and very severe phases of the domain. Meanwhile, there was no significant difference between the total score and scores of psychological and functional domains of severe and very severe stages of COPD [30]. The reason for the contradictory results of this study and other studies might be related to the difference between the asthma and COPD disease and also cultural or religious and differences.

Isobel Stoodley et al. have described asthma as a complex inflammatory disease that causes considerable exhaustion in the patients. Although drug therapy is the dominant method of asthma treatment, there are ever growing evidences that lifestyle interventions can help asthmatic individuals improve their lives. Better nutrition and regular exercises will have ample benefits for the health of the asthmatic patients; however, lengthy research needs to be made to point to the best composition of diet and exercises. In obese asthma patients, weight loss has outspoken benefits. Other lifestyle interventions such as voga and breathing exercises may be helpful for adults suffering from asthma. Massage therapy is more promising for children suffering from

asthma. Despite all the findings, more studies should be conducted to select the best intervention. As a whole, lifestyle interventions are inexpensive treatments that can improve the quality of life and other effects of asthma in both adults and children. Lifestyle interventions should be considered as an essential breakthrough for asthma management [31].

This study was quantitative in design. Future studies are suggested to be qualitative or qualitative-quantitative. Furthermore, since the study data is collected from a single Iranian province, therefore, it is suggested to do a study to collect data from individuals in other provinces and draw an inter-city comparison. The results of this study indicated that the mean score of lifestyle in the asthmatic patients was lower in terms of physical health, exercise and health, psychological, social and environmental health and lower in terms of all aspects of the quality of life.

Acknowledgments

This article is extracted from Master of Science Degree in nursing thesis with the Code A-11-459-3 of Zanjan University of Medical Sciences and Health Services. The authors of this study thank all the respected professors and the Vice-Dean of the Faculty for Research as well as members of the hospitals referred to in this research as well as participants in this study.

Conflict of interest

The authors of this article confirm that there is no conflict of interest with regards to present study.

References

1. Ichionse. M, Sugiura. H, Nagase. H, Yamaguchi. M, Et al. Japanese guidelines for adult asthma 2017. Allergology Int. 2017; 66(2): 163.89.

2. Theresa K, Susan C. Essentials Pediatric

Nursing Children. 2nd ed. United States:

Lippincott Raven. 2016.

3. Özsoy Z, Ekici D. Perception of Nurses Regarding Organizational Justice and Ethical Behavior; Organizational Culture in Hospitals. Hosp Pract Res. 2017; 2(2): 41-46.

4. Ownby Dr. pet dander and difficult to control asthma. the burden of illness. Allergy asthma proc. 2010; 31(5): 381-4.

5. Negahdari S, Ghanbar Zadeh M, Nikbakht M, Tavakol HA. The Effect of 8-Week Aerobic Training on Some Indices of Cardiorespiratory and Exercise Tolerance in Overweight Women's with Chronic Asthma. Jundishapur Sci Med J. 2017; 14(3): 279.93.

6. Walters P, Schofield P, Howard L, Ashworth M, Tylee A. The Relationship between Asthma and Depression in Primary Care Patients: A Historical Cohort and Nested Case Control Study. PLoS ONE. 2011; 6(6): e20750

7. Christian Joachim Apfelbacher. The validity Of Quality Of Life Questionaires in Capturing people's Experience Of Living With Asthma. [dissertation]. 2012.

8. Gruffydd-Jones K, Thomas M, Roman-Rodríguez M, et al. Asthma impacts on workplace productivity in employed patients who are symptomatic despite background therapy: a multinational survey, J Asthma Allergy. 2019, 12: 183-94.

9. Zahran H, Bailey C, P. G. Vital signs: Asthma prevalence, disease characteristics, and self management education.united states, 2001.2009. Morbidity and Mortality Weekly Report. 2011;60(17):547.52.

10. Valizadeh L, Bilan N, Zarei S, Sharifi A. Effect of Education and Controlling Asthma Triggers on Quality of Life among Adolescents with Asthma: A Randomized Clinical Trial. J Mazand Univ Med Sci. 2013; 23(98): 49-57.

11. Amira HA, Amira AA, Mysara MM. Asthma related quality of life in western Saudi subpopulation and its correlation to level of asthma control. Egypt J Chest Dis Tuberc. 2017; 66(1): 11-15.

12. Habib SH, Saha S. Burden of noncommunicable disease: Global overview. Diab Met Syndr Clin Res Rev. 2010; 4(1): 41-47.

13. World Health Organization. The world health report 2002 . Reducing Risks, Promoting Healthy Life. Geneva: World health report, press kit; 2002.

14. National Guide for Asthma Guidelines for Prevention DaT. Tehran: Mehr Tooba; 2015. 58.

15. Samiei Siboni. F, Alimoradi. Z, Sadeghi Z. Impact of corrective life style educational program on controlling stress, anxiety ,and depression in hypertensives. J Birjand Univ Med Sci. 2013; 19(6): 1-9. [In Persian] 16. Bayat F, Shojaeizadeh D, Hosseyni SM, Sadeghi R, Azar T. Effect of Educational Program on Improving Lifestyle in Patients with Type 2 Diabetes. J Res Health Sys. 2013; 8(7): 1235-44. [In Persian]

17. Shiekhy S, Issazadegan A, Khademi A, Hafeznia M. the comparisons of psychopathology dimensions and quality of life(qol) in people with and without asthma. J Urmia Univ Med Sci. 2013; 24 (7): 518-526 URL: <u>http://umj.umsu.ac.ir/article-1-1848-fa.html</u> 18. Megan EJ, Peter GG, Clare EC, Jodi MH, Lisa GW. Lifestyle Risk Factors for Weight Gain in Children with and without Asthma. Children

(Basel). 2017; 4(3): 15. 19. Mahmoudi Rad G, Khaje Mahmoud F, AR. N. Evaluation of the effect of family.centered empowerment model on nutrition and physical activity in patients with asthma. Med Surg Nurs J. 2016; 5(2): 41.48.

20. Bolghan Abadi M, Kimiaee SA, Amir F. The relationship between parents' child rearing styles and their children's quality of life and mental health. Psychology. 2011; 2(3): 230-45.

21. Lali M, Abedi A, Kajbaf MB. Construction and Validation of the Lifestyle Questionnaire (LSQ). Psychol Res. 2012; 15(1): 64.80. [In Persian]

22. Nazem F, Izadi M, Jaliliu M, Keshvarz B. Impact of aerobic exercise and detraining on pulmonary function indexes in obese middle.aged patients with chronic asthma. J Arak Uni Med Sci. 2013; 15(9): 85-93.

23. Saad S. Al Zahrani ,El.Morsy A, El Morsy, Laila Sh.The impact of bronchial asthma on quality of life among affected children and adolescents in Taif city, Saudi Arabia. Life Science Journal. 2014; 11(6). 283-91.

24. Ganjeh P, Mehrabizadeh Honarmand, Davoodi I, Zargar YA. Effectiveness of cognitive.behavioral intervention on quality of life and self.efficacy of boys aged 10 to 15 years with asthma symptoms in Ahvaz city. J Psychol Achieve. 2015; 4(22): 67.88. [In Persian]

25. Stucky BD, Sherbourne CD, Edelen MO, Eberhart NK. Understanding asthma.specific quality of life: moving beyond asthma symptoms and severity. Eur Respir J. 2015; 46: 680-87.

26. Shiekhy S, Issazadegan A, Khademi A, Hafeznia M. The Comparisons Of Psychopathology Dimensions and Quality of life(qol) in people with and without asthma. The Journal of Urmia University of Medical Sciences. 2013; 24(7): 515-26. [In Persian]

27. SalmanYazdi N, Ghasemi I, Salahshourian Fard A, Haghani H.Comparative study on the quality of life amongst asthmatic boys and girls agedbetween 7.17 in Zanjan. Cardiovascular Nurs J; 2012: 1(1): 24-29. [In Persian]

28. Arash M, Shoghi M, Tajvidy M. Assessment Quality of life (QOL) in People with Asthema. Avicenna J Nurs Midwifery care. 2009; 17(12): 13-23.

29. Roohafza HR ,Akochakian SH,Sadeghi M,Fardmoosavi A and etal.Investigating the

Relation between Lifestyle and Chronic Obstractive Pulmonary Disease.Journal .gums.ac.ir.2005,15(59):32.37. [In Persian]

30. Hasanpour N, Attarbashi Moghadam B, Musavi SH, Sami R, Entezari E. The relationship between demographic variables, diseaseseverity and quality of life in chronic obstructive pulmonarydisease patients.Modern Rehabilitation. 2015; 9(3): 19-26. [In Persian]

31. Stoodley I, Williams L, Thompson Ch, Scott H, Wood L. Evidence for lifestyle interventions in asthma. Breathe. 2019; 15: e50-e61.